



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

EXERCISES.

93

FIND the radius of the circle inscribed in the evolute of an ellipse and the ratio in which each point of contact divides the quadrant on which it lies.

[R. H. Graves.]

94

O is the centre of the circumscribed circle of ABC , and D, E, F the middle points of its sides. Show that

$$OD^2 + OE^2 + OF^2 = 2R'(2R' - r'),$$

where R' , r' are the radii of the circumscribed and inscribed circles of the triangle of the feet of the altitudes.

[R. D. Bohannan.]

95

IN exercise 65 let k, l be the lengths of the perpendiculars CD, CE drawn at right angles to CA, CB to meet the base in D, E . Show that

$$\frac{m}{c} = \frac{k}{a} \cdot \frac{l}{b}.$$

[Ormond Stone.]

96

GIVEN on the ground a circular curve of known radius intersecting a given straight line at a given point, and inclined to it at that point at a given angle; it is required to determine the radius of a second circular arc which shall be tangent both to the given curve and to the given line at another given point.

[Calvin Whiteley.]

97

IN the triangle ABC two lines drawn from C trisect the side AB . Given c , C , and the angle φ between the trisecants; to solve the triangle.

[Marcus Baker.]

98

THE eccentric anomalies of three points on an ellipse are p_1, p_2, p_3 . Show that the area of their triangle is

$$\Delta = 2ab \sin \frac{p_2 - p_3}{2} \sin \frac{p_3 - p_1}{2} \sin \frac{p_1 - p_2}{2};$$

the centre of its circumscribing circle

$$x = + \frac{c^2}{a} \cos \frac{p_2 + p_3}{2} \cos \frac{p_3 + p_1}{2} \cos \frac{p_1 + p_2}{2},$$

$$y = - \frac{c^2}{b} \sin \frac{p_2 + p_3}{2} \sin \frac{p_3 + p_1}{2} \sin \frac{p_1 + p_2}{2};$$

and hence show that the centre of curvature of the ellipse at (x, y) is

$$X = +\frac{c^2x^3}{a^4}, \quad Y = -\frac{c^2y^3}{b^4}. \quad [Wm. M. Thornton.]$$

99

FROM a full cask of wine a quantity is taken out at random and the cask filled with water, and then a quantity of the mixture is taken out at random and the cask again filled with water. What is the probability that the cask now contains more wine than water?

[Artemas Martin.]

SELECTED.

100

THE points O, O' defined by the equations in trilinears

$$\begin{aligned} aa : b\beta : c\gamma &= c^2a^2 : a^2b^2 : b^2c^2, \\ aa : b\beta : c\gamma &= a^2b^2 : b^2c^2 : c^2a^2, \end{aligned}$$

are called the Brocard points. The angles $OCA, OAB, OBC, O'BA, O'CB, O'AC$ are called the Brocard angles.

1. Show that the Brocard angles are all equal each to

$$\cot^{-1} [\cot A + \cot B + \cot C].$$

2. Find the equation to the Brocard line OO' .

3. Find the equation to the Brocard circle through O, O' and the centre of the circumscribed circle.

4. Given the base BC and the Brocard angle of a triangle, find the locus of the vertex.

5. Show that the medians bisect the angles between the bisectrices of the angles of ABC and the "symmedian lines"

$$\frac{\alpha}{a} = \frac{\beta}{b} = \frac{\gamma}{c}.$$

6. Show that the Brocard circle contains the symmedian point.

101

EXPRESS in terms of $\sin^{-1}x$ and $\sin^{-1}y$

$$\tan^{-1} \frac{x+y}{\sqrt{(1-x^2)} + \sqrt{(1-y^2)}}.$$

102

FIND the relation connecting x, y, z when

$$\cot^{-1}(x+y+s-xyz) = \cot^{-1}x + \cot^{-1}y + \cot^{-1}z.$$

103

INTO a conical wine glass a spherical ball is dropped. Find the ratio of the concealed surfaces of the ball and the inside of the glass.